Problématique des phtalates et assimilés Où en sommes-nous ?

Point de vue du clinicien

Laurent STORME
Pédiatre Néonatalogiste
CHU & Université de Lille

Déclaration liens d'intérêts

- Aucun

Plastifiants (Phthalates)

Financé par l'ANSM

*Fischer CJ, J Paediatr Child Health. 2013
L Bernard, Sci Total Environ. 2023*

L Bernard, Sci Total Environ. 2023

Quelles conséquences pour les patients ?

- Effets neuro-développementaux
 - Déficit de l'attention
 - Altération cognitive
- Effets perturbateurs endocriniens
 - Effet sur le système hormonal
 - Impact sur la fertilité
- Effets sur le métabolisme
 - Augmentation du risque d'obésité
 - Augmentation de la résistance à l'insuline
- Risques cancérigènes

*Rivollier F, Int J Environ Res Public Health 2019
Raja, Environ Res 2022*

Bisphenol A (BPA)

$I^2 = 0.348$

- 4 cohortes européennes dont EDEN
- N=708 mères enfants
- SDQ test

Mono-n-butyl phthalate (MnBP)

$I^2 < 0.001$

Association entre expositions précoces and devenir

Exposure–outcome associations with a probability of evidence for causation identified up to 2015

	Outcome	Strength of human evidence	Probability of causation, %
Prenatal PBDEs	IQ loss and intellectual disability	Moderate to high	70–100%
Prenatal organophosphate pesticides	IQ loss and intellectual disability	Moderate to high	70–100%
Multiple prenatal exposures	Attention-deficit disorder	Low to moderate	20–69%
Multiple prenatal exposures	Autism spectrum disorder	Low	20–39%
Prenatal DDE	Childhood obesity	Moderate	40–69%
Prenatal BPA	Childhood obesity	Very low to low	20–69%
Adult DEHP	Adult obesity	Low	40–69%
Adult DEHP	Adult diabetes	Low	40–69%
Prenatal DDE	Adult diabetes	Low	20–39%
Prenatal PBDEs	Cryptorchidism	Low	40–69%
Prenatal PBDEs	Testicular cancer	Very low to low	0–19%

Efficacité des interventions ?

**Cosmétiques sans parabène,
triclosan, BP-3 ni phtalates**

Etude avant-après, N=100

RÉSULTATS

Phtalates : c° $\searrow$ 27% MEP (MnBP et MiBP : $\neq$ non significative)

Parabènes : c° $\searrow$ 44% Méthyl et $\searrow$ 45% Propyl. $\nearrow$ c° Ethyl et butyl (mais restent basses)

Triclosan : c° $\searrow$ 36%

Benzophénone-3 : c° $\searrow$ 36 % ($\searrow$ 52% chez les utilisatrices de crème solaire).

Métaanalyses des études interventionnelles visant à réduire l'exposition aux perturbateurs endocriniens

Study author and year	Study design, population, and study size	Intervention active group	Findings
Hagobian, 2017 ⁵⁵	Double-blinded RCT - researchers, lab staff Healthy premenopausal college-aged women with normal BMI (n = 24)	Impact of lifestyle interventions compared to control on urinary BPA concentration. Intervention: Lifestyle interventions include education sessions, BPA-free tupperware/water bottles, makeup, hygiene and feminine products, organic foods packaged in BPA-free glass/cardboard containers. Also included the control intervention of weekly newsletter as below Control: Weekly newsletter on BPA information, healthy eating, and physical activity Duration of Intervention: 3 weeks Summary of findings: Lifestyle Intervention significantly reduced BPA urine concentration	Exposure outcome: Urinary BPA ng/ml Intervention: 0.88, 95% CI 0.42-1.34 Control: 1.37, 95% CI 0.59-2.15
Sathyanarayana, 2012 ⁴⁸	Non-blinded RCT Families - 10 households with 2 or more children aged 4-8 years, 1 parent living 100% in same household as children (n = 40)	Impact of 5-day complete dietary replacement compared to education on urinary phthalates and BPA Intervention: Dietary replacement consistent of fresh/organic food, catered foods prepared without plastics Control: Education alone, conducted via handouts Duration of intervention: 5 days Summary of findings: Dietary intervention caused unexpectedly statistically significant increase in levels of urinary BPA Caused by food contamination with DEHP - high concentrations of DEHP in ground coriander and milk	Exposure outcome: Urinary BPA µg/L Intervention: 1.6, 95% CI 0.9-2.3 Control: 1.4, 95% CI 0.7-2.1

N = 14 études

G Corbet, *Int J Gynecol Obstet* 2022

Etude PREVED – Concentration de perturbateurs endocriniens dans le lait

Table 10. Colostrum biomarkers and exposition in univariate analysis, intervention group with at least two workshop (per protocol analysis) in the PREVED study.

	Control Group (n BPA = 56) (n Parabens = 48)		Intervention Group (n BPA = 91) (n Parabens = 79)		<i>p</i>
	N	%	N	%	
Bisphenol A (BPA)					
Superior to LoD	42	75	71	78	0.67
Inferior to LoD	14	25	20	22	
BPA Mono-Chlorinated (MCBPA)					
Superior to LoD	23	41	44	48	0.39
Inferior to LoD	33	59	47	52	
BPA Di-Chlorinated (DCBPA)					
Superior to LoD	26	46	43	47	0.92
Inferior to LoD	30	54	48	53	
BPA Tri-Chlorinated (TCBPA)					
Superior to LoD	39	70	56	62	0.32
Inferior to LoD	17	30	35	38	
BPA Tetra-Chlorinated (TTBPA)					
Superior to LoD	29	52	46	51	0.88
Inferior to LoD	27	48	45	49	
MethylParaben (MePB)					
Superior to LoD	44	92	68	86	0.34
Inferior to LoD	4	8	11	14	

Etude PREVED – Questionnaires psychologiques

Psychosocial questionnaire					
1-Endocrine-disrupting chemicals knowledge (score/40.5) **	+3.22	[+2.52; +3.93]	+3.57	[+2.99; +4.14]	0.47
2-Perceived ability to avoid chemical exposure (/100) **	+7.09	[+2.33; +11.85]	+5.11	[+1.80; +8.42]	0.50
3-Risk from endocrine-disrupting chemicals (severity score/100) **	+15.61	[+10.64; 20.58]	+21.69	[+18.67; +24.70]	0.03
4-Risk assessment of endocrine-disrupting chemicals (vulnerability score/1400) **	+222.00	[+187.00; +257.10]	+285.10	[+253.60; +316.6]	0.008
Risk perception score (3+4)/100 **	+15.73	[+12.96; +18.51]	+21.03	[+18.99; +23.07]	0.003

Stress périnatal « toxique » (chronique, prolongé, répété)

Est associé à un risque accru de:

- **Pathologies neurodéveloppementales:**

- Déficit d'attention / hyperactivité
- Retard des apprentissages
- Troubles du spectre autistique;

- **Maladies non communicables:**

- Allergie
- Diabète/obésité
- Maladies cardio-vasculaires
- Trouble du comportement: agressivité, contrôle émotionnel, impulsivité;
- Dépression / Anxiété;

Babenko, Neurosci Biobehavior Rev 2015
Boyce, Pediatrics 2021

DOHaD : « Origine Développementale de la Santé »

Facteurs
environnementaux

Fenêtre de vulnérabilité

Santé de l'enfant
et de l'adulte

Pratiques éducatives,
Interactions parents-
enfant,
Langage

Stress « toxique »

Addictions

Activité Physique

Alimentation

Expositions aux Toxiques

*Les premiers
1 000 jours*

Capital Santé & Compétences
psycho-sociales:

- Cognitives
- Sociales
- Emotionnelles

Maladies non communicables:

- Troubles du neuro-développement
- Maladies allergiques / inflammatoires
- Maladies cardiovasculaires
- Maladies métaboliques: obésité, diabète

Impact de l'environnement précoce sur la santé psychique future

Métaanalyse	Odds Ratio (IC 95%)
Toxiques de l'environnement	Estimation entre 1 et 2 (à confirmer)
Sédentarité	1.4 (1.03-1.52)
Obésité	1.4 (1.13-1.71)
Diabète	1.52 (1.23-1.89)
Tabagisme	2.82 (2.38-3.34)
Stress toxique	3.70 (2.62-5.22)
Dépression	4.4 (3.54-5.46)
Addiction	5.62 (4.46-7.07)
Violences intra-familiales	8.10 (5.87-11.18)